

Recommended Adult Immunization Schedule United States, October 2007–September 2008

Note: These recommendations must be read with the footnotes that follow.

Figure 1. Recommended adult immunization schedule, by vaccine and age group

Vaccine ▼	Age group ►	19–49 years	50–64 years	≥65 years
Tetanus, diphtheria, pertussis (Td/Tdap) ^{1,*}		1-dose Td booster every 10 yrs		
		Substitute 1 dose of Tdap for Td		
Human papillomavirus (HPV) ^{2,*}		3 doses (females) (0,2,6 mos)		
Measles, mumps, rubella (MMR) ^{3,*}		1 or 2 doses		1 dose
Varicella ^{4,*}		2 doses (0, 4–8 wks)		
Influenza ^{5,*}		1 dose annually		1 dose annually
Pneumococcal (polysaccharide) ^{6,7}		1–2 doses		1 dose
Hepatitis A ^{8,*}		2 doses (0, 6–12 mos, or 0, 6–18 mos)		
Hepatitis B ^{9,*}		3 doses (0, 1–2, 4–6 mos)		
Meningococcal ^{10,*}		1 or more doses		
Zoster ¹¹				1 dose

*Covered by the Vaccine Injury Compensation Program.

Figure 2. Vaccines that might be indicated for adults based on medical and other indications

Vaccine ▼	Indication ►	Pregnancy	Immuno-compromising conditions (excluding human immunodeficiency virus [HIV]), medications, radiation ¹³	(HIV) infection ^{3,12,13}		Diabetes, heart disease, chronic pulmonary disease, chronic alcoholism	Asplenia ¹² (including elective splenectomy and terminal complement deficiencies)	Chronic liver disease	Kidney failure, end-stage renal disease, recipients of hemodialysis	Healthcare personnel
				CD4+ T lymphocyte count						
				<200 cells/μL	≥200 cells/μL					
Tetanus, diphtheria, pertussis (Td/Tdap) ^{1,*}		1-dose Td booster every 10 yrs								
		Substitute 1 dose of Tdap for Td								
Human papillomavirus (HPV) ^{2,*}		3 doses for females through age 26 years (0, 2, 6 mos)								
Measles, mumps, rubella (MMR) ^{3,*}		Contraindicated								1 or 2 doses
Varicella ^{4,*}		Contraindicated								2 doses (0, 4–8 wks)
Influenza ^{5,*}										1 dose TIV annually
										1 dose TIV or LAIV annually
Pneumococcal (polysaccharide) ^{6,7}										1–2 doses
Hepatitis A ^{8,*}										2 doses (0, 6–12 mos, or 0, 6–18 mos)
Hepatitis B ^{9,*}										3 doses (0, 1–2, 4–6 mos)
Meningococcal ^{10,*}										1 or more doses
Zoster ¹¹		Contraindicated								1 dose

*Covered by the Vaccine Injury Compensation Program.

For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)

Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)

The recommendations in this schedule were approved by the Advisory Committee on Immunization Practices, the American Academy of Family Physicians, the American College of Obstetricians and Gynecologists, and the American College of Physicians.

These schedules indicate the recommended age groups and medical indications for routine administration of currently licensed vaccines for persons aged ≥ 19 years, as of October 1, 2007. Licensed combination vaccines may be used whenever any components of the combination are indicated and when the vaccine's other components are not contraindicated. For detailed recommendations on all vaccines, including those used primarily for travelers or those issued during the year, consult the manufacturers' package inserts and the complete statements from the Advisory Committee on Immunization Practices (www.cdc.gov/vaccines/pubs/acip-list.htm).

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available at www.vaers.hhs.gov or by telephone, 800-822-7967.

Information on how to file a Vaccine Injury Compensation Program claim is available at www.hrsa.gov/vaccinecompensation or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington, D.C. 20005; telephone, 202-357-6400.

Additional information about the vaccines in this schedule, extent of available data, and contraindications for vaccination is also available at www.cdc.gov/vaccines or from the CDC-INFO Contact Center at 800-CDC-INFO (800-232-4636) in English and Spanish, 24 hours a day, 7 days a week.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

Footnotes

Note: Immunization recommendations from ACIP are available at www.cdc.gov/vaccines/pubs/ACIP-list.htm

1. Tetanus, diphtheria, and acellular pertussis (Td/Tdap) vaccination.

Tdap should replace a single dose of Td for adults aged < 65 years who have not previously received a dose of Tdap. Only one of two Tdap products (Adacel[®] [Sanofi Pasteur]) is licensed for use in adults.

Adults with uncertain histories of a complete primary vaccination series with diphtheria and tetanus toxoid-containing vaccines should begin or complete a primary vaccination series. A primary series for adults is 3 doses of tetanus and diphtheria toxoid-containing vaccines; administer the first 2 doses at least 4 weeks apart and the third dose 6–12 months after the second. However, Tdap can substitute for any one of the doses of Td in the 3-dose primary series. The booster dose of tetanus and diphtheria toxoid-containing vaccine should be administered to adults who have completed a primary series and if the last vaccination was received ≥ 10 years previously. Tdap or Td vaccine may be used, as indicated.

If the person is pregnant and received the last Td vaccination ≥ 10 years previously, administer Td during the second or third trimester; if the person received the last Td vaccination in < 10 years, administer Tdap during the immediate postpartum period. A one-time administration of 1 dose of Tdap with an interval as short as 2 years from a previous Td vaccination is recommended for postpartum women, close contacts of infants aged < 12 months, and all healthcare workers with direct patient contact. In certain situations, Td can be deferred during pregnancy and Tdap substituted in the immediate postpartum period, or Tdap can be administered instead of Td to a pregnant woman after an informed discussion with the woman.

Consult the ACIP statement for recommendations for administering Td as prophylaxis in wound management.

2. Human papillomavirus (HPV) vaccination. HPV vaccination is recommended for all females aged ≤ 26 years who have not completed the vaccine series. History of genital warts, abnormal Papanicolaou test, or positive HPV DNA test is not evidence of prior infection with all vaccine HPV types; HPV vaccination is still recommended for these persons.

Ideally, vaccine should be administered before potential exposure to HPV through sexual activity; however, females who are sexually active should still

be vaccinated. Sexually active females who have not been infected with any of the HPV vaccine types receive the full benefit of the vaccination. Vaccination is less beneficial for females who have already been infected with one or more of the HPV vaccine types.

A complete series consists of 3 doses. The second dose should be administered 2 months after the first dose; the third dose should be administered 6 months after the first dose.

Although HPV vaccination is not specifically recommended for females with the medical indications described in Figure 2, "Vaccines that might be indicated for adults based on medical and other indications," it is not a live-virus vaccine and can be administered. However, immune response and vaccine efficacy might be less than in persons who do not have the medical indications described or who are immunocompetent.

3. Measles, mumps, rubella (MMR) vaccination. *Measles component:* adults born before 1957 can be considered immune to measles. Adults born during or after 1957 should receive ≥ 1 dose of MMR unless they have a medical contraindication, documentation of ≥ 1 dose, history of measles based on healthcare provider diagnosis, or laboratory evidence of immunity.

A second dose of MMR is recommended for adults who 1) have been recently exposed to measles or in an outbreak setting; 2) have been previously vaccinated with killed measles vaccine; 3) have been vaccinated with an unknown type of measles vaccine during 1963–1967; 4) are students in postsecondary educational institutions; 5) work in a healthcare facility, or 6) plan to travel internationally.

Mumps component: adults born before 1957 can generally be considered immune to mumps. Adults born during or after 1957 should receive 1 dose of MMR unless they have a medical contraindication, history of mumps based on healthcare provider diagnosis, or laboratory evidence of immunity.

A second dose of MMR is recommended for adults who 1) are in an age group that is affected during a mumps outbreak; 2) are students in postsecondary educational institutions; 3) work in a healthcare facility; or 4) plan to travel internationally. For unvaccinated healthcare workers born before 1957 who do not have other evidence of mumps immunity, consider administering

Footnotes (continued)

Note: Immunization recommendations from ACIP are available at www.cdc.gov/vaccines/pubs/ACIP-list.htm

1 dose on a routine basis and strongly consider administering a second dose during an outbreak.

Rubella component: administer 1 dose of MMR vaccine to women whose rubella vaccination history is unreliable or who lack laboratory evidence of immunity. For women of childbearing age, regardless of birth year, routinely determine rubella immunity and counsel women regarding congenital rubella syndrome. Women who do not have evidence of immunity should receive MMR vaccine on completion or termination of pregnancy and before discharge from the healthcare facility.

4. Varicella vaccination. All adults without evidence of immunity to varicella should receive 2 doses of single-antigen varicella vaccine unless they have a medical contraindication. Special consideration should be given to those who 1) have close contact with persons at high risk for severe disease (e.g., healthcare personnel and family contacts of immunocompromised persons) or 2) are at high risk for exposure or transmission (e.g., teachers; child care employees; residents and staff members of institutional settings, including correctional institutions; college students; military personnel; adolescents and adults living in households with children; nonpregnant women of childbearing age; and international travelers).

Evidence of immunity to varicella in adults includes any of the following: 1) documentation of 2 doses of varicella vaccine at least 4 weeks apart; 2) U.S.-born before 1980 (although for healthcare personnel and pregnant women, birth before 1980 should not be considered evidence of immunity); 3) history of varicella based on diagnosis or verification of varicella by a healthcare provider (for a patient reporting a history of or presenting with an atypical case, a mild case, or both, healthcare providers should seek either an epidemiologic link with a typical varicella case or to a laboratory-confirmed case or evidence of laboratory confirmation, if it was performed at the time of acute disease); 4) history of herpes zoster based on healthcare provider diagnosis; or 5) laboratory evidence of immunity or laboratory confirmation of disease.

Assess pregnant women for evidence of varicella immunity. Women who do not have evidence of immunity should receive the first dose of varicella vaccine upon completion or termination of pregnancy and before discharge from the healthcare facility. The second dose should be administered 4–8 weeks after the first dose.

5. Influenza vaccination: *Medical indications:* chronic disorders of the cardiovascular or pulmonary systems, including asthma; chronic metabolic diseases, including diabetes mellitus, renal or hepatic dysfunction, hemoglobinopathies, or immunosuppression (including immunosuppression caused by medications or human immunodeficiency virus [HIV]); any condition that compromises respiratory function or the handling of respiratory secretions or that can increase the risk of aspiration (e.g., cognitive dysfunction, spinal cord injury, or seizure disorder or other neuromuscular disorder); and pregnancy during the influenza season. No data exist on the risk for severe or complicated influenza disease among persons with asplenia; however, influenza is a risk factor for secondary bacterial infections that can cause severe disease among persons with asplenia.

Occupational indications: healthcare personnel and employees of long-term-care and assisted-living facilities.

Other indications: residents of nursing homes and other long-term-care and assisted-living facilities; persons likely to transmit influenza to persons at high

risk (e.g., in-home household contacts and caregivers of children aged 0–59 months, or persons of all ages with high-risk conditions); and anyone who would like to be vaccinated. Healthy, nonpregnant adults aged ≤ 49 years without high-risk medical conditions who are not contacts of severely immunocompromised persons in special care units can receive either intranasally administered live, attenuated influenza vaccine (FluMist®) or inactivated vaccine. Other persons should receive the inactivated vaccine.

6. Pneumococcal polysaccharide vaccination. *Medical indications:* chronic pulmonary disease (excluding asthma); chronic cardiovascular diseases; diabetes mellitus; chronic liver diseases, including liver disease as a result of alcohol abuse (e.g., cirrhosis); chronic alcoholism; chronic renal failure, or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy [if elective splenectomy is planned, vaccinate at least 2 weeks before surgery]); immunosuppressive conditions; and cochlear implants and cerebrospinal fluid leaks. Vaccinate as close to HIV diagnosis as possible.

Other indications: Alaska Natives and certain American Indian populations and residents of nursing homes or other long-term-care facilities.

7. Revaccination with pneumococcal polysaccharide vaccine. One-time revaccination after 5 years for persons with chronic renal failure or nephrotic syndrome; functional or anatomic asplenia (e.g., sickle cell disease or splenectomy); or immunosuppressive conditions. For persons aged ≥ 65 years, one-time revaccination if they were vaccinated ≥ 5 years previously and were aged < 65 years at the time of primary vaccination.

8. Hepatitis A vaccination. *Medical indications:* persons with chronic liver disease and persons who receive clotting factor concentrates.

Behavioral indications: men who have sex with men and persons who use illegal drugs.

Occupational indications: persons working with hepatitis A virus (HAV)-infected primates or with HAV in a research laboratory setting.

Other indications: persons traveling to or working in countries that have high or intermediate endemicity of hepatitis A (a list of countries is available at www.cdc.gov/travel/content/diseases.aspx) and any person seeking protection from HAV infection.

Single-antigen vaccine formulations should be administered in a 2-dose schedule at either 0 and 6–12 months (Havrix®), or 0 and 6–18 months (Vaqta®). If the combined hepatitis A and hepatitis B vaccine (Twinrix®) is used, administer 3 doses at 0, 1, and 6 months.

9. Hepatitis B vaccination. *Medical indications:* persons with end-stage renal disease, including patients receiving hemodialysis; persons seeking evaluation or treatment for a sexually transmitted disease (STD); persons with HIV infection; and persons with chronic liver disease.

Occupational indications: healthcare personnel and public-safety workers who are exposed to blood or other potentially infectious body fluids.

Behavioral indications: sexually active persons who are not in a long-term, mutually monogamous relationship (e.g., persons with more than one sex partner during the previous 6 months); current or recent injection-drug users; and men who have sex with men.

Other indications: household contacts and sex partners of persons with chronic hepatitis B virus (HBV) infection; clients and staff members of institu-

Footnotes (continued)

Note: Immunization recommendations from ACIP are available at www.cdc.gov/vaccines/pubs/ACIP-list.htm

tions for persons with developmental disabilities; international travelers to countries with high or intermediate prevalence of chronic HBV infection (a list of countries is available at wwwn.cdc.gov/travel/content/diseases.aspx); and any adult seeking protection from HBV infection.

Settings where hepatitis B vaccination is recommended for all adults: STD treatment facilities; HIV testing and treatment facilities; facilities providing drug-abuse treatment and prevention services; healthcare settings targeting services to injection-drug users or men who have sex with men; correctional facilities; end-stage renal disease programs and facilities for chronic hemodialysis patients; and institutions and nonresidential day care facilities for persons with developmental disabilities.

Special formulation indications: for adult patients receiving hemodialysis and other immunocompromised adults, 1 dose of 40 µg/mL (Recombivax HB®) or 2 doses of 20 µg/mL (Engerix-B®), administered simultaneously.

10. Meningococcal vaccination. *Medical indications:* adults with anatomic or functional asplenia or terminal complement component deficiencies.

Other indications: first-year college students living in dormitories; microbiologists who are routinely exposed to isolates of *Neisseria meningitidis*; military recruits; and persons who travel to or live in countries in which meningococcal disease is hyperendemic or epidemic (e.g., the “meningitis belt” of sub-Saharan Africa during the dry season [December–June]), particularly if their contact with local populations will be prolonged. Vaccination is required by the government of Saudi Arabia for all travelers to Mecca during the annual Hajj.

Meningococcal conjugate vaccine is preferred for adults with any of the preceding indications who are aged ≤55 years, although meningococcal polysaccharide vaccine (MPSV4) is an acceptable alternative. Revaccination after 3–5 years might be indicated for adults previously vaccinated with MPSV4 who remain at increased risk for infection (e.g., persons residing in areas in which disease is epidemic).

11. Herpes zoster vaccination. A single dose of zoster vaccine is recommended for adults ≥60 years regardless of whether they report a prior episode of herpes zoster. Persons with chronic medical conditions may be vaccinated unless a contraindication or precaution exists for their condition.

12. Selected conditions for which *Haemophilus influenzae* type b (Hib) vaccine may be used. Hib conjugate vaccines are licensed for children aged 6 weeks–71 months. No efficacy data are available on which to base a recommendation concerning use of Hib vaccine for older children and adults with the chronic conditions associated with an increased risk for Hib disease. However, studies suggest good immunogenicity in patients who have sickle cell disease, leukemia, or HIV infection or who have had splenectomies; administering vaccine to these patients is not contraindicated.

13. Immunocompromising conditions. Inactivated vaccines are generally acceptable (e.g., pneumococcal, meningococcal, influenza [trivalent inactivated influenza vaccine]) and live vaccines generally are avoided in persons with immune deficiencies or immune suppressive conditions. Information on specific conditions is available at www.cdc.gov/vaccines/pubs/acip-list.htm.

The Immunization Action Coalition created this document based on the **Recommended Adult Immunization Schedule, United States, October 2007–September 2008**, published in the *Morbidity and Mortality Weekly Report* on October 19, 2007 (*MMWR* 2007;56:Q1–Q4). This document contains indications for adult immunization based on age (see Figure 1), as well as medical condition and profession (i.e., specific guidance for vaccination of healthcare personnel; see Figure 2). It is also available as a 6-page, 8½” x 11” tri-fold booklet, containing a list of contraindications and precautions for adult immunization, and is laminated and in full color (see ordering information below).

Please note that vaccination recommendations issued by Centers for Disease Control and Prevention (CDC) after the Adult Immunization Schedule’s October 2007 publication date are official even though they are not reflected in this document.

To be sure you have the most current versions of vaccination recommendations, visit the following web pages:

Official ACIP recommendations

www.cdc.gov/vaccines/pubs/acip-list.htm (alphabetical order)

www.immunize.org/acip (chronological order)

Provisional ACIP recommendations

www.cdc.gov/vaccines/recs/provisional/default.htm

www.immunize.org/acip

For other versions of CDC’s adult immunization schedule, go to www.cdc.gov/vaccines/recs/schedules/adult-schedule.htm

To order copies of the laminated, full-color tri-fold booklet described above, visit www.immunize.org/shop, call (651) 647-9009, or email admin@immunize.org.